



[Journals \(/about/journals\)](/about/journals)

[Topics \(/topics\)](/topics)

[Information \(/authors\)](/authors)

[Author Services \(/authors/](/authors)

[toggle desktop layout](#) [cookies](#)

[english](#)

[Initiatives \(/about/initiatives\)](/about/initiatives)

[About \(/about\)](/about)

[Sign In / Sign Up \(/user/login\)](/user/login)

[Submit \(https://susy.mdpi.com/user/manuscripts/upload?journal=fractalfract\)](https://susy.mdpi.com/user/manuscripts/upload?journal=fractalfract)

Search for Articles:

Title / Keyword

Author / Affiliation / Email

Fractal and Fractional (Fractal Fract)

All Article Types

Search

Advanced Search

[Journals \(/about/journals\)](/about/journals) / [Fractal Fract \(/journal/fractalfract\)](/journal/fractalfract) / [Volume 6 \(/2504-3110/6\)](/2504-3110/6) / [Issue 9 \(/2504-3110/6/9\)](/2504-3110/6/9) / [10.3390/fractalfract6090495](https://doi.org/10.3390/fractalfract6090495) /



fractal and fractional **

journal/fractalfract

Submit to this Journal (https://susy.mdpi.com/user/manuscripts/upload?form%5Bjournal_id%5D%3D275)




Review for this Journal (<https://susy.mdpi.com/volunteer/journals/review>)

Propose a Special Issue (/journalproposalsendproposalspecialissue/fractalfract)



Article Menu

Academic Editors

 [./toggle_desktop_layout_cookie](#)  



Hari Mohan Srivastava (https://sciprofiles.com/profile/27527?utm_source=mdpi.com&utm_medium=website&utm_campaign=avatar_name)



Artion Kashuri (https://sciprofiles.com/profile/650368?utm_source=mdpi.com&utm_medium=website&utm_campaign=avatar_name)

[Subscribe SciFeed \(/2504-3110/6/9/495/scifeed_display\)](#)

Recommended Articles

Related Info Link

More by Authors Links

Article Views 1186

Citations 2

Table of Contents

- [Abstract](#)
- [Introduction](#)
- [Preliminaries](#)
- [Weighted Fractional Integral Inequalities](#)
- [Conclusions](#)
- [Author Contributions](#)
- [Funding](#)
- [Institutional Review Board Statement](#)
- [Informed Consent Statement](#)
- [Data Availability Statement](#)
- [Acknowledgments](#)
- [Conflicts of Interest](#)
- [References](#)

IK



Open Access Article

[Order Article Reprints \(/2504-3110/6/9/495/reprints\)](#)



[Share](#)

Certain Weighted Fractional Inequalities via the Caputo–Fabrizio Approach

by **Vaijanath L. Chinchane**¹  (<https://orcid.org/0000-0003-4246-8405>),
Asha B. Nale²  (<https://orcid.org/0000-0003-2793-5992>), **Satish K. Panchal**² and
Christophe Chesneau^{3,*}  (<mailto:christophe.chesneau@unicaen.fr>)

Department of Mathematics, Deogiri Institute of Engineering and Management Studies, Aurangabad 431005, India

Department of Mathematics, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431004, India

Department of Mathematics, University of Caen Normandie, 14000 Caen, France

Author to whom correspondence should be addressed.

Fractal Fract. **2022**, *6*(9), 495; <https://doi.org/10.3390/fractalfract6090495> (<https://doi.org/10.3390/fractalfract6090495>)

Submission received: 10 August 2022 / Revised: 2 September 2022 / Accepted: 3 September 2022 / Published: 5 September 2022

(This article belongs to the Special Issue **Fractional Integral Inequalities and Applications** (/journal/fractalfract/special_issues/fractional_integral_inequalities))

Download 

Versions Notes (/2504-3110/6/9/495/notes)

Abstract

The Caputo–Fabrizio fractional integral operator is one of the important notions of fractional calculus. It is involved in numerous illustrative and practical issues. The main goal of this paper is to investigate weighted fractional integral inequalities using the Caputo–Fabrizio fractional integral operator with non-singular $e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-s)}$, $0 < \delta < 1$. Furthermore, based on a family of n positive functions defined on $[0, \infty)$, we investigate some new extensions of weighted fractional integral inequalities.

Keywords: inequalities (/search?q=inequalities); Caputo–Fabrizio fractional integral operator (/search?q=Caputo%E2%80%93Fabrizio+fractional+integral+operator); weighted fractional inequalities (/search?q=weighted+fractional+inequalities)

1. Introduction

Fractional calculus is the extension of conventional calculus into the non-integer differential and integral orders. It has found significant importance due to its application in various fields of science and engineering, such as life sciences, chemical science, and physical sciences. It naturally appears in

Help

Cite

Discuss in
SciProfiles

[https://](https://sciprofile)

sciprofile

[discussio](https://sciprofile)

[groups/](https://sciprofile)

[public/10](https://sciprofile)

[fractalfra](https://sciprofile)

[utm_sou](https://sciprofile)

Endorse

Comment

MDPI

numerous branches of science and engineering, including the physical, chemical, and life sciences. Numerous mathematicians have spent the last two decades studying fractional integral inequalities and applications using the Riemann–Liouville, Saigo, Hadamard, Marichev–Saigo–Maeda, generalized Katugampola, and generalized k -fractional integral operators, see Refs. [1,2,3,4,5,6,7,8,9,10].

Some recent developments in this field will now be presented. Houas M. looked at certain weighted integral inequalities involving fractional hypergeometric operators in Ref. [11]. Marichev [12] (see also Ref. [10]) introduced the generalization of the hypergeometric fractional integral, including the Saigo operator. Caputo and Fabrizio proposed a novel fractional derivative and application of a new time and spatial fractional derivative with exponential kernels (see Refs. [13,14]). Recently, certain fractional integral inequalities utilizing the Caputo–Fabrizio fractional integral were proposed by Gustava Nchama and et al. in Ref. [15]. In Refs. [16,17], the authors proposed some fractional inequalities for h -convex functions and preinvex functions involving the Caputo–Fabrizio operator. Gürbüz and et al. [18] obtained the Hermite–Hadamard inequality and related inequalities for fractional integrals of Caputo–Fabrizio type. The main motivation is the Caputo–Fabrizio integral and its derivative operator, which are general fractional integral and derivative, respectively (see Refs. [19,20]). Additionally, it has a non-singular kernel, which is a real power that has been transformed into an integral using the Laplace transform. As a result, many problems have an easy time finding an exact solution.

These days, fractional integrals and derivatives play a significant role in modeling a variety of physical phenomena. Due to the singular kernel, the Riemann–Liouville and Caputo fractional derivatives cannot adequately represent several phenomena relating to material heterogeneities. It is the result of Caputo and Fabrizio's suggestion of a novel fractional integral involving the nonsingular kernel $e^{-\left(\frac{1-\delta}{\delta}\right)(x-s)}$, where the parameter δ satisfies $0 < \delta < 1$. Furthermore, the Caputo–Fabrizio fractional integral operator has been increasingly popular among applied scientists' mathematicians as a modeling tool (see Refs. [21,22,23]). In Refs. [15,24], the authors developed fractional integral inequalities using the Caputo–Fabrizio operator. For further information, see Refs. [10,12]. As a concrete physical example, a heat conduction equation with the Caputo–Fabrizio derivative and an investigation of a line segment were utilized to provide the fundamental solutions to the Cauchy and Dirichlet problems in Ref. [25]. The major advantages of the Caputo–Fabrizio integral operator is that it allows the boundary condition of fractional differential equations with Caputo–Fabrizio derivatives to admit the same form as for the differential equations of integer order. A few studies using the Caputo and Caputo–Fabrizio integral operators to solve fractional integral inequalities have been published (see Refs. [15,26,27]). Our goal is to create some novel weighted fractional integral inequalities involving Caputo–Fabrizio fractional integral operators, which are inspired by Refs. [20,22,23,28,29].

The following paragraph describes how the paper has organized. In **Section 2**, basic definitions for the Caputo–Fabrizio fractional derivatives and integrals are provided. In **Section 3**, by utilizing the Caputo–Fabrizio fractional integral operator, we provide weighted fractional integral inequalities. A conclusion is given in **Section 4**.

2. Preliminaries

Here, the Caputo–Fabrizio fractional integral operator is discussed along with some fundamental concepts of fractional calculus.

Definition 1 ([3,15,26]). Let $\delta, a \in \mathbb{R}$ such that $0 < \delta < 1$. The Caputo–Fabrizio fractional derivative of order δ of a function ϕ is defined by

$$\mathcal{J}_{a,\kappa}^{\delta} [\phi(\kappa)] = \frac{1}{1-\delta} \int_a^{\kappa} e^{-\left(\frac{\delta}{1-\delta}\right)(\kappa-s)} \phi'(s) ds. \quad (1)$$

Definition 2 ([3,15,26]). Let $\delta \in \mathbb{R}$ such that $0 < \delta \leq 1$. The Caputo–Fabrizio fractional integral of order δ of a function ϕ is defined by

$$\mathcal{I}_{0,\kappa}^{\delta} [\phi(\kappa)] = \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-s)} \phi(s) ds. \quad (2)$$

For the special case $\delta = 1$, it is reduced to the following integral:

$$\mathcal{I}_{0,\kappa}^1 [\phi(\kappa)] = \int_0^{\kappa} \phi(s) ds.$$

The above definition may be extended to any $\delta > 0$.

With the help of the Caputo–Fabrizio fractional integral operator, various novel inequalities will be demonstrated in this study.

3. Weighted Fractional Integral Inequalities

Using the Caputo–Fabrizio fractional integral operator, we establish a few weighted fractional integral inequalities in this section.

Theorem 1. Let f be a positive continuous function on $[0, \infty)$, and $\varrho > 0$, $\omega \geq \lambda > 0$, such that, for any $\mu, \zeta > 0$,

$$(\zeta^{\varrho} f^{\varrho}(\mu) - \mu^{\varrho} f^{\varrho}(\zeta)) (f^{\omega-\lambda}(\mu) - f^{\omega-\lambda}(\zeta)) \geq 0. \quad (3)$$

In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta > 0$, we have

$$\mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} f^{\omega}(\kappa)] \leq \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\omega}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} f^{\lambda}(\kappa)] \quad (4)$$

Proof. From (3), we have

$$\zeta^{\varrho} f^{\omega-\lambda}(\zeta) f^{\varrho}(\mu) + \mu^{\varrho} f^{\omega-\lambda}(\mu) f^{\varrho}(\zeta) \leq \zeta^{\varrho} f^{\omega+\varrho-\lambda}(\mu) + \mu^{\varrho} f^{\omega+\varrho-\lambda}(\zeta). \quad (5)$$

Then, multiplying both sides of the inequality (5) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) f^{\lambda}(\mu)$, $\mu \in (0, \kappa)$, $\kappa > 0$, and integrating the resultant identity with respect to μ from 0 to κ , we get



$$\begin{aligned}
 & \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \zeta^q f^{\omega-\lambda}(\zeta) w(\mu) f^{q+\lambda}(\mu) d\mu + \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^\omega(\mu) f^q \\
 & \leq \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \zeta^q f^{\omega+q}(\mu) w(\mu) d\mu + \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^{\omega+\lambda}(\mu) f^\lambda(\mu) d\mu
 \end{aligned} \tag{6}$$

consequently,

$$\begin{aligned}
 & \zeta^q f^{\omega-\lambda}(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] + f^q(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] \\
 & \leq \zeta^q \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{\omega+q}(\kappa)] + f^{\omega+\lambda}(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)].
 \end{aligned} \tag{7}$$

Multiplying both sides of (7) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f^\lambda(\zeta)$, $\zeta \in (0, \kappa)$, $\kappa > 0$, then integrating the resulting identity with respect to ζ from 0 to κ , we obtain

$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \\
 & \leq \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{\omega+q}(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa)
 \end{aligned} \tag{8}$$

which brings the proof to a close. \square

The condition (3) is not severe; it can be demonstrated that it is satisfied for any decreasing function f , or any function f such that f and $f(\kappa)/\kappa$ are increasing. We can notice that the inequality (4) becomes an equality for the basic function $f(\kappa) = \kappa$.

We will now present our major finding.

Theorem 2. Let f be a positive continuous function on $[0, \infty)$ and $q > 0$, $\omega \geq \lambda > 0$, which satisfy (3). In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta > 0$, we have

$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \\
 & \leq \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{\omega+q}(\kappa)] + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa)
 \end{aligned} \tag{9}$$

Proof. Let us multiply both sides of (5) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f^\lambda(\zeta)$, ($\zeta \in (0, \kappa)$, $\kappa > 0$), under the circumstances specified in the theorem (this function remaining positive). Then, by integrating the result with respect to ζ from 0 to κ , we obtain

$$\begin{aligned}
 & f^q(\mu) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\zeta) \zeta^q f^\omega(\zeta) d\zeta + \mu^q f^{\omega-\lambda}(\mu) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\zeta) \\
 & \leq f^{\omega+q-\lambda}(\mu) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\zeta) \zeta^q f^\lambda(\zeta) d\zeta + \mu^q \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\zeta) f
 \end{aligned} \tag{10}$$

therefore

$$\begin{aligned}
 & f^q(\mu) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] + \mu^q f^{\omega-\lambda}(\mu) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] \\
 & \leq f^{\omega+q-\lambda}(\mu) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] + \mu^q \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{\omega+q}(\kappa)].
 \end{aligned} \tag{11}$$

Again multiplying both sides of (11) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) f^\lambda(\mu)$, $\mu \in (0, \kappa)$, $\kappa > 0$, then integrating the obtained result with respect to ζ from 0 to κ , we get

$$\begin{aligned} & \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\omega(\kappa)] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) f^{q+\lambda}(\mu) d\mu + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] \\ & \leq \mathcal{I}_{0,\kappa}^\delta [w(\kappa) \kappa^q f^\lambda(\kappa)] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} f^{q+\omega}(\mu) w(\mu) d\mu + \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\omega}(\kappa)] \end{aligned} \quad (12)$$

This completes the proof of the theorem. \square

Theorem 3. Let f and g be positive continuous functions on $[0, \infty)$, and $q > 0$, $\omega \geq \lambda > 0$, such that, for any $\mu, \zeta > 0$, we have

$$(g^q(\zeta) f^q(\mu) - g^q(\mu) f^q(\zeta)) (f^{\omega-\lambda}(\mu) - f^{\omega-\lambda}(\zeta)) \geq 0. \quad (13)$$

In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta > 0$, we get

$$\mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) g^q(\kappa) f^\omega(\kappa)] \leq \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\omega}(\kappa)] \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^\lambda(\kappa)] \quad (14)$$

Proof. From (13), we have

$$g^q(\zeta) f^{\omega-\lambda}(\zeta) f^q(\mu) + g^q(\mu) f^q(\zeta) f^{\omega-\lambda}(\mu) \leq g^q(\zeta) f^{q+\omega-\lambda}(\mu) + g^q(\mu) f^{q+\omega-\lambda}(\zeta) \quad (15)$$


Multiplying both sides of (15) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) f^\lambda(\mu)$, $\mu \in (0, \kappa)$, $\kappa > 0$, then integrating the resulting identity with respect to μ from 0 to κ , we obtain

$$\begin{aligned} & g^q(\zeta) f^{\omega-\lambda}(\zeta) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^{\omega-\lambda}(\mu) [w(\mu) f^{q+\lambda}(\mu)] d\mu \\ & + f^q(\zeta) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^{\omega-\lambda}(\mu) [w(\mu) g^q(\mu) f^\omega(\mu)] d\mu \\ & \leq g^q(\zeta) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^{\omega-\lambda}(\mu) [w(\mu) f^{q+\omega}(\mu)] d\mu \\ & + f^{q+\omega-\lambda}(\zeta) \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \mu^q f^{\omega-\lambda}(\mu) [w(\mu) g^q(\mu) f^\lambda(\mu)] d\mu. \end{aligned} \quad (16)$$

Thus, we establish that

$$\begin{aligned} & g^q(\zeta) f^{\omega-\lambda}(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\lambda}(\kappa)] + f^q(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) g^q(\kappa) f^\omega(\kappa)] \\ & \leq g^q(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) f^{q+\omega}(\kappa)] + f^{q+\omega-\lambda}(\zeta) \mathcal{I}_{0,\kappa}^\delta [w(\kappa) g^q(\kappa) f^\lambda(\kappa)]. \end{aligned} \quad (17)$$

Multiplying both sides of (17) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f^\lambda(\zeta)$, then integrating the resulting inequality with respect to ζ over $(0, \kappa)$, we obtain



$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) g^{\varrho}(\zeta) f^{\omega}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} f^{\varrho+\lambda}(\zeta) w(\zeta) d\zeta \\
& \leq \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\omega+\varrho}(\kappa)] \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} f^{\lambda}(\zeta) w(\zeta) g^{\varrho}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\lambda}(\kappa)] \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f^{\omega+\varrho}(\zeta) d\zeta,
\end{aligned} \tag{18}$$

which implies that

$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [f \\
& \leq \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\omega+\varrho}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [f^{\lambda}(\kappa) w(\kappa) g^{\varrho}(\kappa)] + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\lambda}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta}
\end{aligned} \tag{19}$$

that concludes the proof. \square

It is worth noting that the condition (13) is satisfied for a wide range of function classes. It is satisfied in particular by any functions f and g such that either $f(\kappa)/g(\kappa)$ and $f(\kappa)$ are increasing (simultaneously), or $f(\kappa)/g(\kappa)$ and $f(\kappa)$ are decreasing (simultaneously).

Theorem 4. Let f and g be two positive continuous functions on $[0, \infty)$ and $\varrho > 0$, $\omega \geq \lambda > 0$, which satisfy (13). In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta, \beta > 0$, we have


$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^{\beta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \mathcal{J}_{0,\kappa}^{\beta} [w \\
& \leq \mathcal{J}_{0,\kappa}^{\beta} [w(\kappa) g^{\varrho} f^{\lambda}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\omega+\varrho}(\kappa)] + \mathcal{J}_{0,\kappa}^{\beta} [w(\kappa) f^{\varrho+\omega-\lambda}(\kappa)] \mathcal{J}_{0,\kappa}^{\delta} [w
\end{aligned} \tag{20}$$

Proof. Multiplying the inequality (17) by $\frac{1}{\beta} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) f^{\lambda}(\zeta)$, $\zeta \in (0, \kappa)$, $\kappa > 0$ (this function remains positive under the conditions stated with the theorem), then integrating the obtained result with respect to ζ from 0 to κ , we get


$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \frac{1}{\beta} \int_0^{\kappa} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) g^{\varrho}(\zeta) f^{\omega}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] \frac{1}{\beta} \int_0^{\kappa} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} f^{\varrho+\lambda}(\zeta) w(\zeta) d\zeta \\
& \leq \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) f^{\omega+\varrho}(\kappa)] \frac{1}{\beta} \int_0^{\kappa} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} f^{\lambda}(\zeta) w(\zeta) g^{\varrho}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\beta}(\kappa)] \frac{1}{\beta} \int_0^{\kappa} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) f^{\omega+\varrho}(\zeta) d\zeta,
\end{aligned} \tag{21}$$

which implies that





$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) g^{\varrho}(\kappa) f^{\omega}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] + \mathcal{I}_{0,\kappa}^{\beta} [w(\kappa) f^{\varrho+\lambda}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \\
 & \leq \mathcal{I}_{0,\kappa}^{\beta} [w(\kappa) g^{\varrho} f^{\lambda}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f^{\omega+\varrho}(\kappa)] + \mathcal{I}_{0,\kappa}^{\beta} [w(\kappa) f^{\varrho+\omega}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa)
 \end{aligned} \tag{22}$$

[\(toggle desktop layout cookie\)](#)  

Thus, the proof is completed. \square

In the sequel of the study, we will provide a new extension of weighted fractional integral inequalities based on a family of n positive functions defined on $[0, \infty)$.

Theorem 5. Let $f_i, i = 1, \dots, n$ be n positive continuous functions on $[0, \infty)$ and $\varrho > 0, \omega \geq \lambda_r > 0, r = 1, \dots, n$, such that, for any $\mu, \zeta > 0$,

$$(\zeta^{\varrho} f_r^{\varrho}(\mu) - \mu^{\varrho} f_r^{\varrho}(\zeta)) (f_r^{\omega-\lambda_r}(\mu) - f_r^{\omega-\lambda_r}(\zeta)) \geq 0. \tag{23}$$

In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta > 0$, the following inequality is valid:

$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa)] \\
 & \leq \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} \prod_{i=1}^n f_i^{\lambda_i}(\kappa)] \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f_r^{\omega+\varrho}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa)].
 \end{aligned} \tag{24}$$

Proof. From (23), we have

$$\zeta^{\varrho} f_r^{\omega-\lambda_r}(\zeta) f_r^{\varrho}(\mu) + \mu^{\varrho} f_r^{\varrho}(\zeta) f_r^{\omega-\lambda_r}(\mu) \leq \zeta^{\varrho} f_r^{\omega+\varrho-\lambda_r}(\mu) + \mu^{\varrho} f_r^{\omega+\varrho-\lambda_r}(\zeta). \tag{25}$$

Multiplying both sides of (25) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu)$, then integrating the resulting inequality with respect to μ over $(0, \kappa)$, we obtain

$$\begin{aligned}
 & \zeta^{\varrho} f_r^{\omega-\lambda_r}(\zeta) \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} [w(\mu) f_r^{\varrho}(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu)] d\mu \\
 & + f_r^{\varrho}(\zeta) \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} [w(\mu) \mu^{\varrho} \prod_{i \neq r}^n f_i^{\lambda_i}(\mu)] d\mu \\
 & \leq \zeta^{\varrho} \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} [w(\mu) f_r^{\omega+\varrho}(\mu) \prod_{i \neq r}^n f_i^{\lambda_i}(\mu)] d\mu \\
 & + f_r^{\omega+\varrho-\lambda_r}(\zeta) \frac{1}{\delta} \int_0^{\kappa} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} [w(\mu) \mu^{\varrho} \prod_{i=1}^n f_i^{\lambda_i}(\mu)] d\mu.
 \end{aligned} \tag{26}$$

Consequently,

$$\begin{aligned}
 & \zeta^{\varrho} f_r^{\omega-\lambda_r}(\zeta) \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa)] + f_r^{\varrho}(\zeta) \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa)] \\
 & \leq \sigma^{\varrho} \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) f_r^{\omega+\varrho}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa)] + f_r^{\omega+\varrho-\lambda_r}(\zeta) \mathcal{I}_{0,\kappa}^{\delta} [w(\kappa) \kappa^{\varrho} \prod_{i=1}^n f_i^{\lambda_i}(\kappa)]
 \end{aligned} \tag{27}$$

Again, multiplying the inequality (27) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta)$, $\zeta \in (0, \kappa)$, $\kappa > 0$, this function remaining positive under the conditions stated with the theorem, then integrating the obtained

result with respect to ζ from 0 to κ , we have



$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^e(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) \zeta^e f_r^\omega(\zeta) \prod_{i \neq r}^n f_i^{\lambda_i}(\zeta) d\zeta \\
 & + \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f_r^e(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta) d\zeta \\
 & \leq \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\omega+e}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) \zeta^e \prod_{i=1}^n f_i^{\lambda_i}(\zeta) d\zeta \\
 & + \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\delta} \int_0^\kappa e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) f_r^{\omega+e}(\zeta) \prod_{i \neq r}^n f_i^{\lambda_i}(\zeta) d\zeta,
 \end{aligned} \tag{28}$$

therefore,

$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^e(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[\kappa^e f_r^\omega(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \\
 & + \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^e(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\
 & \leq \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\omega+e}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\
 & + \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\omega+e}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right].
 \end{aligned} \tag{29}$$

This completes the inequality (24). \square

The flexibility of the condition (23) can be commented as the condition (3), but for $f_i, i = 1, \dots, n$ instead of f .

Theorem 6. Let $f_i, i = 1, \dots, n$ be n positive continuous functions on $[0, \infty)$ and $\varrho > 0$, $\omega \geq \lambda_r > 0, r = 1, \dots, n$, which satisfy (23). In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta, \beta > 0$, the following inequality holds:

$$\begin{aligned}
 & \mathcal{I}_{0,\kappa}^\beta \left[w(\kappa) \kappa^e f_r^\omega(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^e(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\
 & + \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e f_r^\omega(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\beta \left[w(\kappa) f_r^e(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\
 & \leq \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\omega+e}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\beta \left[w(\kappa) \kappa^e \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\
 & + \mathcal{I}_{0,\kappa}^\beta \left[w(\kappa) f_r^{\omega+e}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{I}_{0,\kappa}^\delta \left[w(\kappa) \kappa^e \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right].
 \end{aligned} \tag{30}$$

Proof. Multiplying the inequality (27) by $\frac{1}{\beta} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta), \zeta \in (0, \kappa), \kappa > 0$ (under the circumstances outlined in the theorem, this function remains positive), then integrating the result with respect to ζ from 0 to κ , we get



$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^\varrho(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\beta} \int_0^\kappa e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) \zeta^\varrho f_r^\omega(\zeta) \prod_{i \neq r}^n f_i^{\lambda_i}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) \kappa^\varrho \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\beta} \int_0^\kappa e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) f_r^\varrho(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta) d\zeta \\
& \leq \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\varrho+\varrho}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\beta} \int_0^\kappa e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) \zeta^\varrho \prod_{i=1}^n f_i^{\lambda_i}(\zeta) d\zeta \\
& + \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) \kappa^\varrho \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \frac{1}{\beta} \int_0^\kappa e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) f_r^{\varrho+\varrho}(\zeta) \prod_{i \neq r}^n f_i^{\lambda_i}(\zeta) d\zeta,
\end{aligned} \tag{31}$$

which reflects the inequality (30). \square

Theorem 7. Let $f_i, i = 1, \dots, n$ and g be positive continuous functions on $[0, \infty)$, and $\varrho > 0$, $\omega \geq \lambda_r > 0, r = 1, \dots, n$, such that, for any $\mu, \zeta > 0$,

$$(g^\varrho(\zeta) f_r^\varrho(\mu) - g^\varrho(\mu) f_r^\varrho(\zeta)) (f_r^{\omega-\lambda_r}(\mu) - f_r^{\omega-\lambda_r}(\zeta)) \geq 0, \tag{32}$$

In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta > 0$, we have

$$\begin{aligned}
& \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^\varrho(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) g^\varrho(\kappa) f_r^\omega(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \\
& \leq \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) g^\varrho(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\varrho+\varrho}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right].
\end{aligned} \tag{33}$$

Proof. From (32), it is clear that


$$g^\varrho(\zeta) f_r^{\omega-\lambda_r}(\zeta) f_r^\varrho(\mu) + f_r^\varrho(\zeta) g^\varrho(\mu) f_r^{\omega-\lambda_r}(\mu) \leq g^\varrho(\zeta) f_r^{\varrho+\varrho-\lambda_r}(\mu) + f_r^{\varrho+\varrho-\lambda_r}(\zeta) g^\varrho(\mu) \tag{34}$$

Multiplying both sides of (34) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu)$, then integrating the resulting inequality with respect to μ over $(0, \kappa)$, we obtain

$$\begin{aligned}
& g^\varrho(\zeta) f_r^{\omega-\lambda_r}(\zeta) \frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} \left[w(\mu) f_r^\varrho(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu) \right] \\
& + f_r^\varrho(\zeta) \frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} \left[w(\mu) g^\varrho(\mu) f_r^\omega(\mu) \prod_{i \neq r}^n f_i^{\lambda_i}(\mu) \right] \\
& \leq g^\varrho(\zeta) \frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} \left[w(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu) \right] \\
& + f_r^{\varrho+\varrho-\lambda_r}(\zeta) \left[w(\mu) f_r^{\varrho+\varrho}(\mu) g^\varrho(\mu) \prod_{i \neq r}^n f_i^{\lambda_i}(\mu) \right].
\end{aligned} \tag{35}$$

So, we can write

$$\begin{aligned}
& g^\varrho(\zeta) f_r^{\omega-\lambda_r}(\zeta) \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^\varrho(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] + f_r^\varrho(\zeta) \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) g^\varrho(\kappa) f_r^\omega(\kappa) \right] \\
& \leq g^\varrho(\zeta) \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) f_r^{\varrho+\varrho}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] + f_r^{\varrho+\varrho-\lambda_r}(\zeta) \mathcal{J}_{0,\kappa}^\delta \left[w(\kappa) g^\varrho(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right]
\end{aligned} \tag{36}$$

 We multiply both sides of (36) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\zeta)} w(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta)$, and then integrate the resulting inequality with respect to ζ over $(0, \kappa)$, we get the following result:

$$\begin{aligned} & 2 \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \\ & \leq 2 \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right]. \end{aligned} \quad (37)$$

This completes the proof of Theorem 7. \square

Theorem 8. Let $f_i, i = 1, \dots, n$ and g be positive continuous functions on $[0, \infty)$ and $\varrho > 0$, $\omega \geq \lambda_r > 0, r = 1, \dots, n$, such that, for any $\mu, \zeta > 0$,

$$(g^{\varrho}(\zeta) f_r^{\varrho}(\mu) - g^{\varrho}(\mu) f_r^{\varrho}(\zeta)) (f_r^{\omega-\lambda_r}(\mu) - f_r^{\omega-\lambda_r}(\zeta)) \geq 0. \quad (38)$$

In addition, let w be a positive continuous function on $[0, \infty)$. Then, for all $\kappa, \delta, \beta > 0$, the following inequality is valid:

$$\begin{aligned} & \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] + \\ & \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\ & \leq \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\ & + \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right]. \end{aligned} \quad (39)$$

Proof. Multiplying both sides of (34) by $\frac{1}{\beta} e^{-\left(\frac{1-\beta}{\beta}\right)(\kappa-\zeta)} w(\zeta) \prod_{i=1}^n f_i^{\lambda_i}(\zeta)$, then integrating the resulting inequality with respect to ζ over $(0, \kappa)$, we obtain

$$\begin{aligned} & f_r^{\varrho}(\mu) \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] + g^{\varrho}(\kappa) f_r^{\omega-\lambda_r}(\mu) \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\ & \leq f_r^{\varrho+\omega-\lambda_r}(\mu) \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] + g^{\varrho}(\mu) \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \end{aligned} \quad (40)$$

Multiplying both sides of (40) by $\frac{1}{\delta} e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-\mu)} w(\mu) \prod_{i=1}^n f_i^{\lambda_i}(\mu)$, then integrating the resulting inequality with respect to μ over $(0, \kappa)$, we arrive at the following result:

$$\begin{aligned} & \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] + \\ & \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) f_r^{\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] \\ & \leq \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right] + \\ & \mathcal{J}_{0,\kappa}^{\beta} \left[w(\kappa) f_r^{\varrho+\omega}(\kappa) \prod_{i \neq r}^n f_i^{\lambda_i}(\kappa) \right] \mathcal{J}_{0,\kappa}^{\delta} \left[w(\kappa) g^{\varrho}(\kappa) \prod_{i=1}^n f_i^{\lambda_i}(\kappa) \right]. \end{aligned} \quad (41)$$

Thus, the proof is completed. \square

4. Conclusions



Numerous mathematicians continue to be interested in the use of various fractional operators in the study of integral inequalities. Refs. [6,7,30] investigated weighed fractional integral inequalities employing, respectively, the Hadamard, Marichev–Saigo–Maeda, and generalized Katugampola fractional integral operators. While using the Caputo–Fabrizio fractional integral operators, we looked at several new fractional integral inequalities in this research. Due to its nonsingular kernel, i.e., $e^{-\left(\frac{1-\delta}{\delta}\right)(\kappa-s)}$ with $0 < \delta < 1$, it reveals to be a more powerful fractional operator. By taking into account the Caputo–Fabrizio fractional integral operator, we investigated the newly weighted fractional integral inequalities in this study. Using the inequalities discussed in this paper, future research should be able to prove the existence and uniqueness of a number of ordinary differential equations as well as initial and boundary value problems using Caputo–Fabrizio fractional operators. It is also possible that the inequalities found in this study provide access to various relationships between fractals and Caputo–Fabrizio fractionals.

Author Contributions

Conceptualization, V.L.C., A.B.N., S.K.P. and C.C. methodology, V.L.C., A.B.N., S.K.P. and C.C.; validation, V.L.C., A.B.N., S.K.P. and C.C.; investigation, V.L.C., A.B.N., S.K.P. and C.C.; writing—original draft preparation, V.L.C., A.B.N., S.K.P. and C.C.; writing—review and editing, V.L.C., A.B.N., S.K.P. and C.C. All authors have read and agreed to the published version of the manuscript.

Funding

This research received no external funding.

Institutional Review Board Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

Not applicable.

Acknowledgments

We would like to thank the six reviewers and the associate editor for the constructive and thorough comments on the paper.

Conflicts of Interest

The authors declare that they have no competing interests.



1. Anastassiou, G.A. *Fractional Differentiation Inequalities*; Springer Publishing Company, Incorporated: New York, NY, USA, 2009. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fractional+Differentiation+Inequalities&author=Anastassiou,+G.A.&publication_year=2009)]
2. Baleanu, D.; Purohit, S.D.; Agarwal, P. On fractional integral inequalities involving Hypergeometric operators. *Chin. J. Math.* **2014**, *2014*, 609476. [Google Scholar (https://scholar.google.com/scholar_lookup?title=On+fractional+integral+inequalities+involving+Hypergeometric+operators&author=Baleanu,+D.&author=Purohit,+S.D.&author=Agarwal,+P.&publication_year=2014&journal=Chin.+J.+Math.&volume=2014&pages=609476&doi=10.1155/2014/609476)] [CrossRef (<https://doi.org/10.1155/2014/609476>)]
3. Nale, A.B.; Chinchane, V.L.; Panchal, S.K.; Chesneau, C. Pólya-Szegő integral inequalities using the Caputo-Fabrizio approach. *Axioms* **2022**, *11*, 79. [Google Scholar (https://scholar.google.com/scholar_lookup?title=P%C3%B3lya-Szeg%C3%B6+integral+inequalities+using+the+Caputo-Fabrizio+approach&author=Nale,+A.B.&author=Chinchane,+V.L.&author=Panchal,+S.K.&author=Chesneau,+C.&publication_year=2022&journal=Axioms&volume=11&pages=79&doi=10.3390/axioms11020079)] [CrossRef (<https://doi.org/10.3390/axioms11020079>)]
4. Kilbas, A.A.; Srivasatava, H.M.; Trujillo, J.J. *Theory and Application of Fractional Differential Equation*; North-Holland Mathematiccal Studies Vol; Elsevier Science Publishers: Amsterdam, The Netherlands; London, UK; New York, NY, USA, 2006. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Theory+and+Application+of+Fractional+Differential+Equation&author=Kilbas,+A.A.&author=Srivasatava,+H.M.&author=Trujillo,+J.J.&publication_year=2006)]
5. Nale, A.B.; Panchal, S.K.; Chinchane, V.L. Certain fractional integral inequalities using generalized Katugampola fractional integral operator. *Malaya J. Math.* **2020**, *3*, 791–797. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Certain+fractional+integral+inequalities+using+generalized+Katugampola+fractional+integral+operator&author=Nale,+A.B.&author=Panchal,+S.K.&author=Chinchane,+V.L.&publication_year=2020&journal=Malaya+J.+Math.&volume=3&pages=791%E2%80%97797&doi=10.26637/MJM0803/0013)] [CrossRef (<https://doi.org/10.26637/MJM0803/0013>)]





- Nale, A.B.; Panchal, S.K.; Chinchane, V.L. Weighted fractional inequalities using Marichev-Saigo-Maeda fractional integral operator. *J. Korean Soc. Ind. Appl. Math.* **2021**, *25*, 39–53. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Weighted+fractional+inequalities+using+Marichev-Saigo-Maeda+fractional+integral+operator&author=Nale,+A.B.&author=Panchal,+S.K.&author=Chinchane,+V.L.&publication_year=2021&journal=J.+Korean+Soc.+Ind.+Appl.+Math.&volume=25&pages=39%E2%80%9353)]
7. Nale, A.B.; Panchal, S.K.; Chinchane, V.L. On weighted fractional inequalities using Hadamard fractional integral operator. *Palest. J. Math.* **2021**, *10*, 614–624. [Google Scholar (https://scholar.google.com/scholar_lookup?title=On+weighted+fractional+inequalities+using+Hadamard+fractional+integral+operator&author=Nale,+A.B.&author=Panchal,+S.K.&author=Chinchane,+V.L.&publication_year=2021&journal=Palest.+J.+Math.&volume=10&pages=614%E2%80%93624)]
8. Podlubny, I. *Fractional Differential Equations*; Academic Press: Cambridge, MA, USA; New York, NY, USA, 1999. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fractional+Differential+Equations&author=Podlubny,+I.&publication_year=1999)]
9. Saigo, M. A remark on integral operators involving the Gauss hypergeometric functions. *Math. Rep. Kyushu Univ.* **1978**, *11*, 135–143. [Google Scholar (https://scholar.google.com/scholar_lookup?title=A+remark+on+integral+operators+involving+the+Gauss+hypergeometric+functions&author=Saigo,+M.&publication_year=1978&journal=Math.+Rep.+Kyushu+Univ.&volume=11&pages=135%E2%80%93143)]
10. Somko, S.G.; Kilbas, A.A.; Marichev, O.I. *Fractional Integral and Derivative Theory and Application*; Gordon and Breach: Yverdon, Switzerland, 1993. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fractional+Integral+and+Derivative+Theory+and+Application&author=Somko,+S.G.&author=Kilbas,+A.A.&author=Marichev,+O.I.&publication_year=1993)]
11. Houas, M. Certain weighted integral inequalities involving the fractional hypergeometric operators. *Sci. Ser. Math. Sci.* **2016**, *27*, 87–97. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Certain+weighted+integral+inequalities+involving+the+fractional+hypergeometric+operators&author=Houas,+M.&publication_year=2016&journal=Sci.+Ser.+Math.+Sci.&volume=27&pages=87%E2%80%9397)]
12. Marichev, O.I. Volterra equation of Mellin convolution type with a Horn function in the kernel. *Izv. AN BSSR Ser. Fiz. Mat. Nauk* **1974**, *1*, 128–129. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Volterra+equation+of+Mellin+convolution+type+with+a+Horn+function+in+the+kernel&author=Marichev,+O.I.&publication_year=1974&journal=Izv.+AN+BSSR+Ser.+Fiz.+Mat.+Nauk&volume=1&pages=128%E2%80%93129)]





- Caputo, M.; Fabrizio, M. A new Definition of Fractional Derivative without singular kernel. *Progr. Fract. Differ. Appl.* **2015**, *1*, 73–85. [Google Scholar (https://scholar.google.com/scholar_lookup?title=A+new+Defination+of+Fractional+Derivative+withou+singular+kernel&author=Caputo,+M.&author=Fabrizio,+M.&publication_year=2015&journal=Progr.+Fract.+Differ.+Appl.&volume=1&pages=73%E2%80%9385)]
14. Caputo, M.; Fabrizio, M. Applications of new time and spatial fractional derivative with exponential kernels. *Progr. Fract. Differ. Appl.* **2016**, *2*, 7–8. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Applications+of+new+time+and+spatial+fractional+derivative+with+exponential+kernels&author=Caputo,+M.&author=Fabrizio,+M.&publication_year=2016&journal=Progr.+Fract.+Differ.+Appl.&volume=2&pages=7%E2%80%938&doi=10.18576/pfda/020101)] [CrossRef (<https://doi.org/10.18576/pfda/020101>)]
15. Nchama, G.A.M.; Mecias, A.L.; Richard, M.R. The Caputo-Fabrizio fractional integral to generate some new inequalities. *Inf. Sci. Lett.* **2019**, *2*, 73–80. [Google Scholar (https://scholar.google.com/scholar_lookup?title=The+Caputo-Fabrizio+fractional+integral+to+generate+some+new+inequalities&author=Nchama,+G.A.M.&author=Mecias,+A.L.&author=Richard,+M.R.&publication_year=2019&journal=Inf.+Sci.+Lett.&volume=2&pages=73%E2%80%9380)]
16. Chen, L.; Zhang, J.; Saleem, M.S.; Ahmed, I.; Waheed, S.; Pan, L. Fractional integral inequalities for h-convex functions via Caputo-Fabrizio operator. *AIMS Math.* **2021**, *6*, 6377–6389. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fractional+integral+inequalities+for+h-convex+functions+via+Caputo-Fabrizio+operator&author=Chen,+L.&author=Zhang,+J.&author=Saleem,+M.S.&author=Ahmed,+I.&author=Waheed,+S.&author=Pan,+L.&publication_year=2021&journal=AIMS+Math.&volume=6&pages=6377%E2%80%936389&doi=10.3934/math.2021374)] [CrossRef (<https://doi.org/10.3934/math.2021374>)]
17. Tariq, M.; Ahmad, H.; Shaikh, A.G.; Sahoo, S.K.; Khedher, K.M.; Gia, T.N. New fractional integral inequalities for preinvex functions involving Caputo-Fabrizio operator. *AIMS Math.* **2022**, *7*, 3440–3455. [Google Scholar (https://scholar.google.com/scholar_lookup?title=New+fractional+integral+inequalities+for+preinvex+functions+involving+Caputo-Fabrizio+operator&author=Tariq,+M.&author=Ahmad,+H.&author=Shaikh,+A.G.&author=Sahoo,+S.K.&author=Khedher,+K.M.&author=Gia,+T.N.&publication_year=2022&journal=AIMS+Math.&volume=7&pages=3440%E2%80%933455&doi=10.3934/math.2022191)] [CrossRef (<https://doi.org/10.3934/math.2022191>)]






- Gürbüz, M.; Akdemir, A.O.; Rashid, S.; Set, E. Hermite-Hadamard inequality for fractional integrals of Caputo-Fabrizio type and related inequalities. *J. Inequal. Appl.* **2020**, *172*, 1–17. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Hermite-Hadamard+inequality+for+fractional+integrals+of+Caputo-Fabrizio+type+and+related+inequalities&author=G%C3%BCrb%C3%BCz,+M.&author=Akdemir,+A.O.&author=Rashid,+S.&author=Set,+E.&publication_year=2020&journal=J.+Inequal.+Appl.&volume=172&pages=1%E2%80%9317&doi=10.1186/s13660-020-02438-1)] [CrossRef (<https://doi.org/10.1186/s13660-020-02438-1>)]
19. Abdeljawad, T.; Baleanu, D. On fractional derivatives with exponential kernel and their discrete versions. *Rep. Math. Phys.* **2017**, *1*, 11–27. [Google Scholar ([https://scholar.google.com/scholar_lookup?title=On+fractional+derivatives+with+exponential+kernel+and+their+discrete+versions&author=Abdeljawad,+T.&author=Baleanu,+D.&publication_year=2017&journal=Rep.+Math.+Phys.&volume=1&pages=11%E2%80%9327&doi=10.1016/S0034-4877\(17\)30059-9](https://scholar.google.com/scholar_lookup?title=On+fractional+derivatives+with+exponential+kernel+and+their+discrete+versions&author=Abdeljawad,+T.&author=Baleanu,+D.&publication_year=2017&journal=Rep.+Math.+Phys.&volume=1&pages=11%E2%80%9327&doi=10.1016/S0034-4877(17)30059-9))] [CrossRef ([https://doi.org/10.1016/S0034-4877\(17\)30059-9](https://doi.org/10.1016/S0034-4877(17)30059-9))]
20. Abdeljawad, T. Fractional operators with exponential kernels and a Lyapunov type inequality. *Adv. Differ. Equ.* **2017**, *2017*, 1–11. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fractional+operators+with+exponential+kernels+and+a+Lyapunov+type+inequality&author=Abdeljawad,+T.&publication_year=2017&journal=Adv.+Differ.+Equ.&volume=2017&pages=1%E2%80%9311&doi=10.1186/s13662-017-1285-0)] [CrossRef (<https://doi.org/10.1186/s13662-017-1285-0>)]
21. Atanackovic, T.M.; Janev, M.; Pilipovic, S. Wave equation in fractional Zener-type viscoelastic media involving Caputo-Fabrizio fractional derivatives. *Meccanica* **2019**, *54*, 155–167. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Wave+equation+in+fractional+Zener-type+viscoelastic+media+involving+Caputo-Fabrizio+fractional+derivatives&author=Atanackovic,+T.M.&author=Janev,+M.&author=Pilipovic,+S.&publication_year=2019&journal=Meccanica&volume=54&pages=155%E2%80%93167&doi=10.1007/s11012-018-0920-5)] [CrossRef (<https://doi.org/10.1007/s11012-018-0920-5>)]
22. Dokuyucu, M.A. A fractional order alcoholism model via Caputo-Fabrizio derivative. *Aims Math.* **2020**, *5*, 781–797. [Google Scholar (https://scholar.google.com/scholar_lookup?title=A+fractional+order+alcoholism+model+via+Caputo-Fabrizio+derivative&author=Dokuyucu,+M.A.&publication_year=2020&journal=Aims+Math.&volume=5&pages=781%E2%80%93797&doi=10.3934/math.2020053)] [CrossRef (<https://doi.org/10.3934/math.2020053>)]





- Alshabanat, A.; Jleli, M.; Kumar, S.; Samet, B. Generalization of Caputo-Fabrizio Fractional Derivative and Applications to Electrical Circuits. *Front. Phys.* **2020**, *8*, 7–11. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Generalization+of+Caputo-Fabrizio+Fractional+Derivative+and+Applications+to+Electrical+Circuits&author=Alshabanat,+A.&author=Jleli,+M.&author=Kumar,+S.&author=Samet,+B.&publication_year=2020&journal=Front.+Phys.&volume=8&pages=7%E2%80%9311&doi=10.3389/fphy.2020.00064)] [CrossRef (<https://doi.org/10.3389/fphy.2020.00064>)]
24. Wang, X.; Saleem, M.S.; Aslam, K.N.; Wu, X.; Zhou, T. On Caputo-Fabrizio fractional integral inequalities of Hermite-Hadamard type for modified-convex functions. *J. Math.* **2020**, *2020*, 8829140. [Google Scholar (https://scholar.google.com/scholar_lookup?title=On+Caputo-Fabrizio+fractional+integral+inequalities+of+Hermite-Hadamard+type+for+modified-convex+functions&author=Wang,+X.&author=Saleem,+M.S.&author=Aslam,+K.N.&author=Wu,+X.&author=Zhou,+T.&publication_year=2020&journal=J.+Math.&volume=2020&pages=8829140&doi=10.1155/2020/8829140)] [CrossRef (<https://doi.org/10.1155/2020/8829140>)]
25. Hristov, J.; Bennacer, R. *Heat Conduction: Methods, Applications and Research*; Nova Science Publishers: Hauppauge, NY, USA, 2019. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Heat+Conduction:+Methods,+Applications+and+Research&author=Hristov,+J.&author=Bennacer,+R.&publication_year=2019)]
26. Chinchane, V.L.; Nale, A.B.; Panchal, S.K.; Chesneau, C. On some fractional integral inequalities involving Caputo-Fabrizio integral operator. *Axioms* **2021**, *10*, 255. [Google Scholar (https://scholar.google.com/scholar_lookup?title=On+some+fractional+integral+inequalities+involving+Caputo-Fabrizio+integral+operator&author=Chinchane,+V.L.&author=Nale,+A.B.&author=Panchal,+S.K.&author=Chesneau,+C.&publication_year=2021&journal=Axioms&volume=10&pages=255&doi=10.3390/axioms10040255)] [CrossRef (<https://doi.org/10.3390/axioms10040255>)]
27. Nale, A.B.; Panchal, S.K.; Chinchane, V.L. Grüss-type fractional inequality via Caputo-Fabrizio integral operator. *RGMI Res. Rep. Coll.* **2021**, *24*, 1–15. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Gr%C3%BCss-type+fractional+inequality+via+Caputo-Fabrizio+integral+operator&author=Nale,+A.B.&author=Panchal,+S.K.&author=Chinchane,+V.L.&publication_year=2021&journal=RGMI+Res.+Rep.+Coll.&volume=24&pages=1%E2%80%9315)]




-  Al-Refai, M.; Jarrah, A.M. Fundamental results on weighted Caputo-Fabrizio fractional derivative. *Chaos Solitons Fractals* **2019**, *126*, 7–11. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Fundamental+results+on+weighted+Caputo-Fabrizio+fractional+derivative&author=Al-Refai,+M.&author=Jarrah,+A.M.&publication_year=2019&journal=Chaos+Solitons+Fractals&volume=126&pages=7%E2%80%9311&doi=10.1016/j.chaos.2019.05.035)] [CrossRef (<https://doi.org/10.1016/j.chaos.2019.05.035>)]
29. Fadugba, S.E. Development of a new numerical scheme for the solution of exponential growth and decay models. *Open J. Math. Sci.* **2021**, *4*, 18–26. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Development+of+a+new+numerical+scheme+for+the+solution+of+exponential+growth+and+decay+models&author=Fadugba,+S.E.&publication_year=2021&journal=Open+J.+Math.+Sci.&volume=4&pages=18%E2%80%9326&doi=10.30538/oms2021.0141)] [CrossRef (<https://doi.org/10.30538/oms2021.0141>)]
30. Panchal, S.K.; Chinchane, V.L.; Nale, A.B. On weighted fractional inequalities using generalized Katugampola fractional integral operator. *Fract. Differ. Calc.* **2020**, *10*, 255–266. [Google Scholar (https://scholar.google.com/scholar_lookup?title=On+weighted+fractional+inequalities+using+generalized+Katugampola+fractional+integral+operator&author=Panchal,+S.K.&author=Chinchane,+V.L.&author=Nale,+A.B.&publication_year=2020&journal=Fract.+Differ.+Calc.&volume=10&pages=255%E2%80%93266&doi=10.7153/fdc-2020-10-16)] [CrossRef (<https://doi.org/10.7153/fdc-2020-10-16>)]

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>) (<https://creativecommons.org/licenses/by/4.0/>).

Share and Cite

 (mailto:?)
 &subject=From%20MDPI%3A%20%22Certain%20Weighted%20Fractional%20Inequalities%20via%20the%20Caputo%26ndash%3BFabrizio%20Approach"&body=https://www.mdpi.com/1813484%3A%0A%0ACertain%20Weighted%20Fractional%20Inequalities%20via%20the%20Caputo%26ndash%3BFabrizio%20Approach%0A%0AAbstract%3A%20The%20Caputo%26ndash%3BFabrizio%20fractional%20integral%20operator%20is%20one%20of%20the%20important%20notions%20of%20fractional%20calculus.%20%20is%20involved%20in%20numerous%20illustrative%20and%20practical%20issues.

The main goal of this paper is to investigate weighted fractional integral inequalities using the Caputo–Fabrizio fractional integral operator with non-singular $e^{-\delta|t-\tau|}$ kernel.

Furthermore, based on a family of non-positive functions defined on $[0, \infty)$, we investigate some new extensions of weighted fractional integral inequalities. (https://twitter.com/intent/tweet?text=Certain+Weighted+Fractional+Inequalities+via+the+Caputo%E2%80%93Fabrizio+Approach&hashtags=mdpifracfract&url=https%3A%2F%2Fwww.mdpi.com%2F1813484&via=Fractal_Fract)

(http://www.linkedin.com/shareArticle?mini=true&url=https%3A%2F%2Fwww.mdpi.com%2F1813484&title=Certain%20Weighted%20Fractional%20Inequalities%20via%20the%20Caputo%E2%80%93Fabrizio%20Approach%26source%3Dhttps%3A%2F%2Fwww.mdpi.com%26summary%3DThe%20Caputo%E2%80%93Fabrizio%20fractional%20integral%20operator%20is%20one%20of%20the%20important%20notions%20of%20fractional%20calculus.

It is involved in numerous illustrative and practical issues. The main goal of this paper is to investigate weighted fractional integral inequalities. (https://www.facebook.com/sharer.php?u=https://www.mdpi.com/1813484) (http://www.reddit.com/submit?url=https://www.mdpi.com/1813484) (http://www.mendeley.com/import/?url=https://www.mdpi.com/1813484)

MDPI and ACS Style

Chinchane, V.L.; Nale, A.B.; Panchal, S.K.; Chesneau, C. Certain Weighted Fractional Inequalities via the Caputo–Fabrizio Approach. *Fractal Fract.* **2022**, *6*, 495. https://doi.org/10.3390/fractalfract6090495

AMA Style


Chinchane VL, Nale AB, Panchal SK, Chesneau C. Certain Weighted Fractional Inequalities via the Caputo–Fabrizio Approach. *Fractal and Fractional*. 2022; 6(9):495. https://doi.org/10.3390/fractalfract6090495

Chicago/Turabian Style

Chinchane, Vaijanath L., Asha B. Nale, Satish K. Panchal, and Christophe Chesneau. 2022. "Certain Weighted Fractional Inequalities via the Caputo–Fabrizio Approach" *Fractal and Fractional* 6, no. 9: 495. https://doi.org/10.3390/fractalfract6090495

Article Metrics





For more information on the journal statistics, click [here \(/journal/fractalfract/stats\)](/journal/fractalfract/stats).

ⓘ Multiple requests from the same IP address are counted as one view.

[Fractal Fract. \(/journal/fractalfract\)](/journal/fractalfract), EISSN 2504-3110, Published by MDPI

[RSS \(/rss/journal/fractalfract\)](/rss/journal/fractalfract) **[Content Alert \(/journal/fractalfract/toc-alert\)](/journal/fractalfract/toc-alert)**

Further Information

[Article Processing Charges \(/apc\)](/apc)

[Pay an Invoice \(/about/payment\)](/about/payment)

[Open Access Policy \(/openaccess\)](/openaccess)

[Contact MDPI \(/about/contact\)](/about/contact)

[Jobs at MDPI \(https://careers.mdpi.com\)](https://careers.mdpi.com)

Guidelines

[For Authors \(/authors\)](/authors)

[For Reviewers \(/reviewers\)](/reviewers)

[For Editors \(/editors\)](/editors)

[For Librarians \(/librarians\)](/librarians)

[For Publishers \(/publishing_services\)](/publishing_services)

[For Societies \(/societies\)](/societies)

[Conference Organizers \(/conference_organizers\)](/conference_organizers)

[MDPI Initiatives](#)

[Sciforum \(https://sciforum.net\)](https://sciforum.net)

MDPI
[MDPI Books \(https://www.mdpi.com/books\)](https://www.mdpi.com/books)

[Preprints.org \(https://www.preprints.org\)](https://www.preprints.org)

[\(\(toggle_desktop_layout_cookie\)\)](#) 🔍 ☰

[Scilit \(https://www.scilit.net\)](https://www.scilit.net)

[SciProfiles \(https://sciprofiles.com?utm_source=mpdi.com&utm_medium=bottom_menu&utm_campaign=initiative\)](https://sciprofiles.com?utm_source=mpdi.com&utm_medium=bottom_menu&utm_campaign=initiative)

[Encyclopedia \(https://encyclopedia.pub\)](https://encyclopedia.pub)

[JAMS \(https://jams.pub\)](https://jams.pub)

[Proceedings Series \(/about/proceedings\)](/about/proceedings)

Follow MDPI

[LinkedIn \(https://www.linkedin.com/company/mdpi\)](https://www.linkedin.com/company/mdpi)

[Facebook \(https://www.facebook.com/MDPIOpenAccessPublishing\)](https://www.facebook.com/MDPIOpenAccessPublishing)

[Twitter \(https://twitter.com/MDPIOpenAccess\)](https://twitter.com/MDPIOpenAccess)



Subscribe to receive issue release notifications and newsletters from MDPI journals

Select options ▼

Enter your email address...

Subscribe

© 1996-2024 MDPI (Basel, Switzerland) unless otherwise stated

[Disclaimer](#)

[Terms and Conditions \(/about/terms-and-conditions\)](/about/terms-and-conditions)

[Privacy Policy \(/about/privacy\)](/about/privacy)

