




Influence of light wavelengths, light intensity, temperature, and pH on biosynthesis of extracellular and intracellular pigment and biomass of *Pseudomonasaeruginosa NR1*

[Naziya N. Mohammad Aziz Rehman](#) , [Prashant P. Dixit](#)  

[Show more](#) 

 Outline |  Share  Cite

<https://doi.org/10.1016/j.jksus.2019.01.004> 

[Get rights and content](#) 

Under a Creative Commons [license](#) 

open access

Abstract

The purpose of the present study was to check whether the light is an inducing agent or not for the intracellular and extracellular pigment production by *Pseudomonas aeruginosa NR1*? Is the produced pigment a photoprotective in nature? The aim of the present research study was to evaluate the influence of different light wavelengths, light intensity, temperature, and pH on biomass and intracellular and extracellular pigment production. A strain of *Pseudomonas* that was thermoalkalophilic and produces red non-diffusible intracellular and green-diffusible extracellular pigment was isolated, until now no one has reported this type of strain. For determining the effect of different light wavelengths, we introduced and designed a novel 'selective light wavelength method', which gave us better results than previously used glass paper wrapping method. Extracellular pigment production was found maximum under darkness, red and blue light wavelengths. Same results were obtained in glass paper wrapping method, but the concentration of pigment produced by bacteria was greater in our method which

suggests us that the 'selective light wavelength method' is better (for pigment yield) between these methods.

The outcome of this study refutes the photo protective role of extracellular pigment, claimed by many researchers. Red and light wavelength was found to suppress the intracellular pigment production. Extracellular pigment production by *Pseudomonas aeruginosa* NR1 was maximum in darkness with 4.042 ± 0.03 OD/mL and minimum in green light (Glass paper plates (GPP), 2.151 ± 0.24 OD/mL; Selective lamp plates (SLP), 2.232 ± 0.28 OD/mL). Maximum intracellular pigment (SLP, 4.672 ± 0.15 OD/mL; GPP, 4.395 ± 0.03 OD/mL) and biomass concentrations (Selective lamp method- 830 ± 5.7 μ g/0.01 L; Glass paper method, 810 ± 2.88 μ g/0.01 L) were achieved under yellow light wavelength. Minimum intracellular pigment yield (GPP, 2.124 ± 0.01 OD/mL; SLP, 2.252 ± 0.0 OD/mL) and biomass yield (Glass paper method, 363.3 ± 3.33 μ g/0.01 L; selective lamp method- 385 ± 2.88 μ g/0.01 L) was recorded under red light wavelength. Direct relationship was observed between biomass production and intracellular pigment production. Maximum extracellular (3.548 ± 2.2 OD/mL) and intracellular pigment (3.856 ± 0.02 OD/mL) and biomass concentration (800 ± 1.44 μ g/100 mL) were observed when pH was 7. The intracellular pigment concentration (3.698 ± 0.01 OD/ml) and biomass production (656 ± 3.05 μ g/ml) were maximal at a temperature of 30 °C. The maximum concentration of extracellular pigment was recorded at 35 °C (3.924 ± 0.04 OD/ml).



Keywords

Pseudomonas aeruginosa NR1; Light wavelength; Light intensity; Biomass; Intracellular; Extracellular pigment concentration

1. Introduction

Environmental factors such as temperature, light etc., are important for regulating the developmental and physiological processes for humans, animals and microbes. We are well familiar with the fact that humans and animals sense and behave accordingly to temperature, light and other many environmental factors. Microbes also can sense, respond and behave accordingly in different environmental conditions and therefore the ability to sense and respond to light in microbes is considered evolutionary conserved mechanism (Arden et al., 1966). For microbial growth and secondary metabolite production, environmental factors are very important. There is a high influence of

medium composition and the operating conditions e.g., temperature, pH, availability and type of light wavelength, degree of aeration etc., on microbial cell growth, development, reproduction and metabolite production. This influence has always been a big hindrance in optimizing the biotechnological process (Chen et al., 2015). Optimization of bacterial and fungal pigment production has become an important aspect now days because many microbial pigments have been reported to have medicinal properties such as anticancer (Ferreira et al., 2004, Kodach et al., 2006), anti protozoan (Matz et al., 2004), antiviral (Sánchez et al., 2006), antibacterial (Nakamura et al., 2003), antioxidant (Konzen et al., 2006), anti-hepatotoxic (Gallardo-Casas et al., 2010), antidiabetic (Soni et al., 2009), or anti-inflammatory (Gitte et al., 2015) activity.

There is a very strong influence of light and temperature on bacterial and fungal pigment production. In this context, *Serratia marcescens* can produce prodigiosin only at low temperature (28 °C), but not at 37 °C (Xu et al., 2014, Nora et al., 2017). Light-dependent accumulation of carotenoid pigments has been studied in *Neurospora* (Rau and Mitzka-Schnabel, 1985). In fungi, many types of photo receptors have been identified (Corrochano, 2007, Herrera-Estrella and Horwitz, 2007). Light has been an inducing agent for pigment (carotenoid) production for several species of non photosynthetic representatives of the genera *Mycobacterium*, *Flavobacterium dehydrogenans*, *Myxococcus*, *Micrococcus roseus* (Yasumasa et al., 1975). All these reports suggest us that there is a huge influence of these factors on pigment production, and they also regulate the pigment production mechanism. As bacteria also sense and respond to light, there is a possibility that bacteria could also possess photo receptors like fungi, which should be detected. Considering all these facts and possibilities, in present research study, the effect of presence or absence of light (total darkness), type of light wavelength, light intensity, temperature and pH on *Pseudomonas aeruginosa NR1* growth, and on intracellular and extracellular pigment and biomass production was investigated.

2. Materials and methods

2.1. Materials

All the glassware and Petri dishes used in experimentation were purchased from Scott Duran manufacturer (DURAN®, Germany). The glassware consists of good quality glass with clear, colourless properties, having negligibly low absorption spectra in the range about 310 to 2200 nm. Petri dishes used in experiments could transmit light of all wavelengths by more than 90%. Philips 20Watts lamps (Philips India Ltd.) were used as a light source for experimentation. All other chemicals used in experimentation were of analytical grade, and were procured from Merck, India. Light intensity was controlled in

both experimental setups by using the same 20Watt lamps, and by keeping all plates and flasks equidistant (15 cm) from the light source.

2.2. Isolation of pigment producing bacterium


The pigment producing bacterium was isolated from dead Rhinoceros beetle. The insect was collected aseptically and was brought into the laboratory. The feathers of the insect were cut and were washed with sterile distilled water. This water was serially diluted up to a factor of 10^{-6} with sterile distilled water. Diluted samples were used for the isolation of pigmented bacteria. A loopful of diluted sample from 10^{-6} tube was withdrawn with Nichrome wire loop, and was streaked on sterile nutrient agar (Hi-Media, Mumbai) plates. Plates were incubated at 37°C in incubator for 48 h. A colony with intense and bright pigment producing ability was selected for the further research studies. The selected bacterium was able to produce dark red intracellular ([Fig. 3](#)) and green extracellular pigments ([Fig. 4](#)).

```
GGAAACATTTATGCGACTTCCCCAGTCATGAATCACTCCGTGGTAACCGTCCTCCTT
GCGGTTAGACTAGCTACTTCTGGAGCAACCCACTCCCATGGTGTGACGGGCGGTGTGT
ACAAGGCCCGGGAACGTATTCACCGTGACATTCGTGATTCACGATTACTAGCGATTCCG
ACTTCACGCAGTCGAGTTGCAGACTGCGATCCGGACTACGATCGGTTTTATGGGATTA
GCTCCACCTCGCGGCTTGGCAACCCCTTTGTACCGACCATTTGTAGCACGTGTGTAGCC
TGGCCGTAAGGGCCATGATGACTTGACGTCAATCCCACTTCCTCCGGTTTGTACCCG
GCAGTCTCCTTAGAGTGCCCAACCGAGGTGCTGGTAACTAAGGACAAGGGTTGCGCT
CGTTACGGGACTTAACCCAACATCTCACGACACGAGCTGACGACAGCCATGCAGCAC
CTGTGTCTGAGTTCCTCGAAGGCA
CCGAGGTAAGGTTCTTGGGTTTC
```

[Download](#) : [Download high-res image \(530KB\)](#)

[Download](#) : [Download full-size image](#)


Fig. 1. . 16S rRNA of *Pseudomonas aeruginosa* NR1.



[Download : Download high-res image \(71KB\)](#)

[Download : Download full-size image](#)


Fig. 2. . Phylogenetic tree of *P. aeruginosa* NR1.



[Download : Download high-res image \(94KB\)](#)

[Download : Download full-size image](#)

Fig. 3. . Intracellular pigment produced by *P. aeruginosa* NR1.



[Download](#) : [Download high-res image \(67KB\)](#)

[Download](#) : [Download full-size image](#)

Fig. 4. . Extracellular pigment produced by *P. aeruginosa* NR1.

2.3. Identification of isolated pigmented bacterium

The isolated bacterium was identified by morphological, biochemical characteristics, and by the 16S rRNA gene sequencing method. The 16S rRNA gene was amplified by using following 16S universal primers.

16S Forward primer: 5' AGA GTT TGA TCC TGG CTC AG 3'

16S Reverse primer: 5' AAG GAG GTG ATC CAG CCG CA 3'

2.4. Effect of different light wavelengths on pigment and biomass production

Effect of different light wavelengths on pigment and biomass production was evaluated by using two methods. One was glass paper wrapping method, which has been used by researchers ([Soumya et al., 2014](#), [Babitha et al., 2008](#)) to deduce effect of light wavelengths on microbial pigment and biomass production; the second was by using the selective lamp light wavelength method, which is a novel technique and for the first time introduced by our research group. First method was based on the principle that the color glass paper allows only selective wavelength of light to pass through it. If red colored glass paper is used, then only red color light wavelength will pass through the paper. In the second method, only selective colored light wavelength was provided to flasks and plates, which were kept in controlled confined environment. Colored lamps (Philips, 20W) were used for providing selective wavelengths.

2.5. Experimental set up for method one (glass wrapping method)

For this method, five different color glass papers were used. The papers used in this research study were green, red, yellow, brown, blue in color (Fig. 5). Each glass paper was wrapped around the one flask and a plate and these wrapped plates and flasks were kept around the illumination source, 15 cm away from the light source (Fig. 6). One flask and one plate were directly kept under illumination without wrapping to check effect of direct light on pigment and biomass production. To detect effect of darkness on pigment and biomass production, one flask and one plate were kept in total darkness. For intracellular pigment production, the organism was grown on sterile nutrient agar plates; for extracellular pigment production, as it was diffusible only in liquid medium, the organism was grown in sterile nutrient broth flasks. Flasks and plates were inoculated with the organism with equal volumes (2 mL) of 24 h old bacterial culture and were incubated in a box that had a single illumination source provided centrally.

[Download](#) : [Download high-res image \(110KB\)](#)

[Download](#) : [Download full-size image](#)

Fig. 5. . Glass papers wrapped to plates and flasks.

[Download](#) : [Download high-res image \(140KB\)](#)

[Download](#) : [Download full-size image](#)

Fig. 6. Wrapped flasks and plated under light source.

2.6. Experimental set up for method two (selective wavelength method)

We designed 'a selective light wavelength' experiment for evaluating the effect of light on pigment and biomass production. For this experiment, a wooden box was constructed. This box was equally subdivided in nine chambers by partitioning the box with wooden blocks (Fig. 7). The partition was done in such a manner that no light rays from one chamber could pass to another chamber. An electric holder plate was designed with 10 lamp holders (Fig. 8). This plate was designed in such a way that in each chamber a holder socket was present in a central position. At every holder socket, one colored selective wavelength lamp was fixed. The box was closed, so that no external light was able to enter the box. Red, orange, blue, green, yellow, white, pink lamps and one more lamp with multiple light wavelengths that exerts red, blue, green, yellow, white light

wavelength one by one repeatedly were used for the study. In one chamber no lamp was fixed, so that total darkness is achieved for incubation in absence of light.

[Download : Download high-res image \(87KB\)](#)

[Download : Download full-size image](#)

Fig. 7. . Equal chambers made in box.

[Download : Download high-res image \(71KB\)](#)

[Download : Download full-size image](#)

Fig. 8. . Electric plate constructed for the experiment.

2.7. Effect of temperature on pigment and biomass production.

To evaluate effect of temperature on pigment and biomass production, sterile nutrient broth flasks and nutrient agar plates were prepared. For each temperature, one flask and one plate were investigated. All plates and flasks were inoculated with equal amounts of bacterial culture and kept at different temperatures such as 25 °C, 35 °C, 45 °C, 55 °C, 65 °C. After 2 days of incubation, the amount of pigment and biomass was estimated.

2.8. Effect of pH on pigment and biomass production.

To check the effect of pH, one nutrient broth flask and one nutrient agar plate with pH 3, 4, 5, 6, 7, 8, 9, 10 were prepared and inoculated with equal amounts of bacterial culture and were kept for incubation for 2 days at 30 °C. Amount of pigment and biomass was estimated by using spectrophotometric analysis (Elico UV–Visible double beam spectrophotometer).

2.9. Effect of light intensity on pigment production

For detecting the effect of increased light intensity on extracellular and intracellular pigment production, and to get information regarding role of extracellular pigment in photoprotection, this study was undertaken. To study the light intensity effect on pigment production, the plates and flasks producing intracellular and extracellular pigment, respectively, were kept and incubated under increasing light intensity environment by using 15 W, 20 W, 25 W, 30 W, 40 W, 50 W white light lamps.

2.10. Inoculation

2 mL of inoculum medium was used for all flasks. Twenty-four hours old broth of *P. aeruginosa* NR1 was used for inoculation. Plates and flasks were inoculated with desired bacterial culture and all were kept for incubation at their places for 72 h at 35 °C. The room temperature was maintained at 35 °C with the help of room heater.

2.11. Intracellular and extracellular pigment estimation

Extracellular pigment concentration was estimated by measuring absorption of pigmented solution at its λ_{max} , i.e., 382 nm. Broth was centrifuged with relative centrifugal force of $3600 \times g$ for 20 min by using REMI research centrifuge (REMI, India). The supernatant was filtered and cleared, colored filtrate was obtained. Equal amount of Chloroform was used for the complete extraction of extracellular pigment from the aqueous broth. The colored chloroform was used for spectrophotometric analysis (Elico UV-Visible double beam spectrophotometer). For estimating intracellular pigment concentration, the intracellular pigment was extracted by using acetone. 10 mL of acetone was added in a Petri plate with well grown pigmented colonies. The pigment slowly started to dissolve in solvent. When colonies turned colorless, the acetone was removed and collected in a test tube. 5 mL of acetone was again added to extract the remaining traces of pigment completely from colonies. The red colored acetone was filtered with Whatmann filter paper GF/C disc (47 mm). The absorption maxima were determined for the red pigment, and the concentration of pigment in each plate was estimated by measuring the absorption in spectrophotometer (Elico UV-Visible double beam spectrophotometer). Acetone was used as blank in the analysis.

2.12. Biomass estimation

The bacterial biomass was estimated as dry weight by the conventional method described by [Li and Mira \(2010\)](#) with some minor modification. Broth was centrifuged with relative centrifugal force of $3600 \times g$ for 20 min by using REMI research centrifuge (REMI, Centrifuge, India) for 20 min. The pellet was washed with distilled water and was mixed in 5 mL distilled water. The empty weight (in gram) of the evaporating dish was measured and recorded (initial weight, IW). The bacterial mass was kept in evaporating dish. The dish was kept in oven at 100 °C until all water was evaporated, and only bacterial mass remained in dish. After complete drying, the final weight (in gram) of the dish with the dried bacterial mass was recorded (Final weight, FW). The actual weight of bacterial cells was determined by subtracting the initial weight from the final weight.

$$\text{Dry weight} = \text{FW} - \text{IW}$$

where,

FW – final dish weight (with dried bacterial mass)

IW – initial dish (empty dish weight)

2.13. Statistical analysis

Each experiment was repeated at least three times. Numerical data are presented as mean \pm SD. All the results were analyzed by means of one-way ANOVA. All statistical analyses were performed by using SPSS 11.5 software. $P < 0.05$, $P < 0.01$ was considered statistically significant.

3. Result and discussion

3.1. Isolation and identification of pigmented bacterium-

The red colored bacterium was identified as *P. aeruginosa* NR1 by 16srRNA gene sequencing and considering it's morphological and biochemical characteristics. The bacterium was able to catabolize many sugars including glucose, galactose, sucrose, maltose, fructose, mannose, dextrose, xylose, but was found unable to catabolize rhamnose, lactose, arabinose, and mannitol (Table 1). Moreover, the bacterium was oxidase and catalase positive. The 16S rRNA gene sequence and phylogenic tree is shown in Fig. 1, Fig. 2 respectively.

Table 1. Biochemical characteristics of *P. aeruginosa* NR1 (VP- Voges–Proskauer).

| Glucose Fermentation | Sucrose fermentation | Fructose fermentation | Lactose fermentation | Arabinose Fermentation | Galactose fermentation | Dextrose fermentation | | |
|----------------------|----------------------|-----------------------|----------------------|------------------------|------------------------|-----------------------|------------------|-------------------|
| Positive | Positive | Positive | Negative | Negative | Positive | Positive | | |
| Xylose fermentation | Maltose fermentation | Indole test | Methyl red test | VP Test | Citrate Test | Amylase enzyme | Chitinase enzyme | Gelatinase Enzyme |
| Positive | Positive | Negative | Negative | Positive | Positive | Positive | Positive | Negative |

3.2. Effect of light wavelengths, light intensity, temperature and pH on pigment and biomass production.

[Propst and Lubin \(1979\)](#) have also reported the light mediated changes in *P. aeruginosa* cultures. In their research study, they found that cultures of *Pseudomonas aeruginosa* PA0 grown under uninterrupted broad-spectrum light showed pigmentation different from dark-grown cultures; dark-grown bacteria produced pigments, which resulted in blue-purple colored agar, while light-grown organisms produced red colored plates. We didn't observe complete change in pigment color, but a slight change in color shade was

observed when our bacterium was kept under light and total darkness. In our experiment, higher concentration of pigment was obtained for the flasks and plates of 'selective wavelengths lamp light method' than for the 'glass paper wrapped' flasks and plates (Table 2). This higher concentration of pigment in selective wavelengths method shows its superiority and efficacy over the glass paper wrapping method. For intracellular pigment production, yellow light wavelength was found most suitable in our experiment. Under yellow light wavelength, maximum intracellular pigment production was recorded (SLP- 4.672 ± 0.1 OD/ml; GPP- 4.395 ± 0.03 OD/ml), followed by green (SLP - 3.816 ± 0.02 OD/ml; GPP - 3.611 ± 0.07 OD/ml), orange (SLP - 3.500 ± 0.03 OD/ml), and pink (SLP - 2.913 ± 0.04 OD/ml). Minimum intracellular pigment production was observed under red (GPP - 2.124 ± 0.01 OD/ml; SLP- 2.252 ± 0.02 OD/ml), blue (GPP- 2.308 ± 0.01 OD/ml; SLP - 2.412 ± 0.02 OD/ml) and white light wavelengths (SLP - 2.890 ± 0.03 OD/ml). Our light experiments revealed that under blue and white light, intracellular pigment production decreases, which is in accordance with the findings by Nobutaka et al., 2004, Chen et al., 2016. As a difference between our result and those by Nobutaka et al. (2004) results, red light negatively influenced the pigment formation in our experiments, while in Nobutaka et al. (2004) red light didn't affect the pigment and biomass concentration. Our result that under blue light the intracellular pigment production decreases, was similar with the result by De Lucca et al., (2012), as they reported that the blue light effectively inhibits bacterial and fungal growth, whereas Miyake et al. (2005) in a research study reported opposite observation that under blue light pigment production was maximum by *Monascus ruber* N. Darkness was not that much effective for the organism for intracellular pigment production. On the other hand, extracellular pigment was produced maximum under darkness (4.042 ± 0.13 OD/mL). This result is in accordance with Babitha et al., 2008, Soumya et al., 2014, and Velmurugan et al., 2010, who also reported that darkness was beneficial for the pigment production by their organism. Our this result is significant as this observation refutes the photo protective role of our pigment as proposed by Dieringer et al., 1977, Yong and Lee, 1991, Salih et al., 2000, and Seagle et al., (2005). Using red light wavelength (Selective lamp flask 3.714 ± 0.2 OD/mL; Glass paper flask, 3.317 ± 0.07 OD/mL) and blue wavelength (Selective lamp flask 3.638 ± 0.19 ; Glass paper flask, 3.590 ± 0.02), extracellular pigment production was also pronounced. The lowest extracellular pigment concentration was obtained under green light wavelength (GPP- 2.151 OD/mL; SLP- 2.232 OD/mL). The biomass was produced maximum under yellow (Selective lamp flask 830 ± 5.7 μ g/0.01 L; Glass paper flask, 810 ± 2.88 μ g/0.01 L) and green light (Selective lamp flask 731 ± 4.4 μ g/0.01 L; Glass paper flask, 708 ± 4.4 μ g/0.01 L) wavelengths. This means that the biomass and intracellular pigment was produced at the maximum concentration under yellow and green light. This result was expected since the concentration of intracellular pigment should be directly proportional to the concentration of biomass; however, in a research published by Alejandro et al. (2011), the opposite observation was reported

namely no direct relationship between biomass concentration and pigment production. Pigment production by the organism was not completely inhibited under any particular light wavelength.

Table 2. Effect of light wavelengths on intracellular, extracellular pigment and biomass production by *P. aeruginosa* NR1 by glass paper wrapping method and selective light wavelength method. All results are expressed as mean \pm standard deviation.

| Concentration | Type of light | | | | | | |
|---------------------------------|------------------------|-------------------------|-----------------------------|------------------------------|------------------|-------------------------------|---------------------------|
| | Red Paper (620–750) | Blue Paper (450–495) | Green Paper (495–570) | Yellow Paper (570–590) | Brown paper | Red Lamp Bulb (620–750) | Blue Bulb (450–495) |
| Intracellular pigment 540 nm | 2.124 \pm 0.01 | 2.308 \pm 0.0 | 3.611 \pm 0.07 | 4.395 \pm 0.03 | 2.721 \pm 0.02 | 2.2352 \pm | 2.124 \pm 0.01 |
| Extracellular pigment 382 nm | 3.317 \pm 0.007 | 3.590 \pm 0.02 | 2.151 \pm 0.24 | 3.206 \pm 0.32 | 3.296 \pm 0.25 | 3.714 \pm 0.21 | 3.317 \pm 0.007 |
| Biomass(μ g/10 mL) | 363.3 \pm 3.33 | 413 \pm 3.33 | 708 \pm 4.40 | 810 \pm 2.88 | 650 \pm 2.88 | 385 \pm 2.88 | 413 \pm 3.33 |

Table 3. Effect of temperature on intracellular, extracellular pigment and biomass production by *P. aeruginosa* NR1. All results are expressed as mean \pm standard deviation. Symbols-+++ indicates very good growth; ++ indicates good growth; + indicates very less growth.

| Temperature ($^{\circ}$ C) \rightarrow | 25 | 30 | 35 | 40 |
|--|----------------------------------|------------------------------|------------------------------|-----------------------------|
| Factor \downarrow | | | | |
| Intracellular pigment/color 540 nm | 4.539 \pm 0.04/orange | 3.698 \pm 0.01/dark Red | 4.433 \pm 0.02/dark red | 3.414 \pm 0.2/lig pink |
| Extracellular 382 nm pigment/absorbance/color | 2.448 \pm 0.02/yellow green | 3.257 \pm 0.07/green | 3.924 \pm 0.04/green | 3.230 \pm 0.01/G |
| Growth | Yes/++ | Yes/+++ | Yes/+++ | Yes/++ |
| Biomass(microgram/ml) | 618.3 \pm 4.4 | 656 \pm 3.05 | 592.3 \pm 1.452 | 551.6 \pm 6.0 |

Table 4. Effect of pH on intracellular, extracellular pigment and biomass production by *P. aeruginosa* NR1. All results are expressed as mean \pm standard deviation. Symbols-+++ indicates very good growth; ++ indicates good growth; + indicates very less growth.

| pH \rightarrow at 33 °C Factor \downarrow | pH 3 | pH 4 | pH 5 | pH 6 | pH 7 | pH 8 |
|--|------|------|---|------------------------------|-------------------------------|-----------------------|
| Intracellular pigment/color 540 nm | No | No | 2.548 \pm 0.35/light pink | 3.753 \pm 2.14/dark red | 3.856 \pm 0.02/dark red | 3.524 \pm red |
| Extracellular pigment | No | No | 2.861 \pm 1.86 light Yellowish green | 3.048 \pm 0.75 green | 3.548 \pm 2.2 Dark Green | 3.564 \pm Dark G |
| Growth | No | No | Yes/+ | Yes/++ | Yes/+++ | +++/Gr |
| Biomass(microgram/100 mL) | - | - | 500 \pm 0.50 | 650 \pm 1.12 | 800 \pm 1.44 | 800 \pm 0 |

In our research study, we used two methods to detect the effect of light wavelengths on pigment production. Both methods, i.e., glass paper wrapping method and selective wavelength lamp method, gave good results, but more satisfactory and more efficient results were obtained in method two, i.e., selective lamp light wavelength method; as the pigment concentrations were higher under selective light lamps than the glass paper wrapped flasks and plates. In different light wavelengths, we observed some differences in color shade of extracellular pigment. This might be due to the variation in composition of pigment as suggested by [Velmurugan et al. \(2010\)](#). Our result and conclusion is also in agreement with [Miyake et al. \(2005\)](#), who stated that pigment concentration changes with the variation in light wavelength. When bacterial plates and flasks were incubated under increasing light intensity, no considerable changes in pigment concentration were observed ([Table 5](#)). These results also refute the role of our pigment in photoprotection. The effect of temperature on extracellular and intracellular pigment and biomass production is shown in [Table 3](#). The maximum extracellular (3.924 \pm 0.04 OD/mL) and intracellular pigment concentration (3.698 \pm 0.01 OD/mL) was recorded when temperature was 35 °C and 30 °C, respectively. Maximum biomass concentration (656 \pm 3.05 μ g/0.01 L) was obtained when temperature was 30 °C. Above 45 °C, neither intra- nor extracellular pigment production was observed. Growth was observed up to 55 °C, but not at higher temperature. This shows that this bacterium is a moderate thermophile. Maximum intracellular pigment (3.856 \pm 0.02 OD/mL), extracellular pigment (3.548 \pm 2.2 OD/ml) and biomass concentration (800 μ g/0.01 L) was recorded at pH 7. Above pH 10 and below pH 5, no pigment production was recorded. The bacterial growth was observed up to pH 10, which means the isolated bacterium is an alkalophilic strain. [Table 4](#) shows the effect of pH on extracellular, intracellular pigment and biomass

production by the bacterium. It is apparent from this research study that the pigment of isolated bacterium doesn't have any role in photoprotection. This fact raises many questions in mind like why does the bacterium produce the pigment? Many bacterial and fungal pigments have been well recognized for their virulent activities (Liu and Nizet, 2009, Hall et al., 2016, Ortiz-Castro et al., 2014, Carbonell et al., 2000). As we had isolated the pigment producing bacterium from dead insect, we may speculate that the pigment of the bacterium might have acted as virulent factor for the insect and it might have served a reason for insect's death. Another role of the pigment (considering its source of isolation) might be in degradation of cellular components (like chitin) of insect for making energy for bacterial cellular processes. This presumption seems more convincing after considering the ability of the bacterium to produce chitinase enzyme (the isolated bacterium was able to produce chitin in colossal amount). The pigment might be having roles other than these two hypothesized roles. The pigment may help the bacterium in competition with other microbes for getting needed nutrients for its growth and survival or it could have a role in bacterial resistance to naturally produced antimicrobial compounds by other microbes or plants. The pigment might be helpful for the bacterium to cope up with harsh and extreme conditions like high temperature, high pressure, high salinity and cold. Many researchers have reported the role of pigments in salt tolerance by haloarchae (Rebecca et al., 2016, Oren, 2010).

Table 5. Effect of light intensity on pigment production by *Pseudomonas aeruginosa* NR1. All results are expressed as mean \pm standard deviation.

| Light intensity | 15 Watt | 20 Watt | 30 Watt | 40 Watt | 50 Watt | Darkness |
|---------------------------------------|-----------------|------------------|------------------|------------------|-----------------|------------------|
| Extracellular pigment OD at 382 nm | 2.431 \pm 0.2 | 2.315 \pm 0.33 | 2.442 \pm 0.03 | 2.492 \pm 0.2 | 2.496 \pm 1.1 | 4.154 \pm 0.15 |
| Intracellular pigment OD at 540 nm | 2.875 \pm 1.3 | 2.877 \pm 0.33 | 2.814 \pm 0.1 | 2.770 \pm 0.05 | 2.814 \pm 4.1 | 3.296 \pm 0.35 |

4. Conclusion

From this study, we conclude that we have isolated a thermoalkalophilic strain of *P. aeruginosa* that produces red non-diffusible intracellular and green diffusible extracellular pigments. We also infer that there is an influence of light wavelength, pH and temperature on pigment production and biomass production of *P. aeruginosa* NR1. However, temperature displays the major influence factor, because slight difference in temperature completely inhibited the pigment production. Light intensity doesn't affect considerably on pigment production by the bacterium. From this study, we infer that there is a strong effect of different light wavelengths on pigment production by *P.*

aeruginosa NR1 as pigment production was doubled under suitable light wavelength. Biomass production is also influenced by pH and temperature. For pH and temperature, this was expected as it is well known fact that bacterial growth is influenced by pH and temperature. However, our study demonstrates that bacterial growth and therefore biomass production is also influenced by light wavelength. We conclude that darkness, red and blue light are suitable conditions for maximum extracellular pigment production, while yellow and green light is suitable for maximum biomass and intracellular pigment production by *P. aeruginosa NR1*. We also conclude that our designed selective wavelength lamp model for light experiments is a more suitable technique than the paper wrapping method. Our model is also advantageous over paper wrapping method as, once designed, it is easy to handle and also gives maximum pigment yield.

Conflict of interest

The authors declare that they have no conflict of interest.

Statement of human rights/animal welfare

This article does not contain any studies with human participants or animals performed by any of the authors.

Informed consent

“Informed consent was obtained from all individual participants included in the study.”

Acknowledgment

The authors are thankful to DST INSPIRE, New Delhi, India for providing fellowship to first author (IF150419) of this manuscript.

[Recommended articles](#)

References

- [Alejandro et al., 2011](#) M. Alejandro, P. Catalina, C.M. Julio, M. Gabriela, Noé A. Cristóbal
Red pigment production by *Penicillium purpurogenum* GH2 is influenced by pH and temperature
J. Zhejiang Univ. Sci. B, 12 (12) (2011), pp. 961-968, [10.1631/jzus.B1100039](#) [↗](#)
[Google Scholar](#) [↗](#)

[Arden et al., 1966](#) G.B. Arden, H. Ikeda, I.M. Siegal

New components of the mammalian receptor potential and their relation to visual photochemistry

Vision Res., 6 (1966), p. 373, [10.1016/0042-6989\(66\)90047-2](#) ↗

 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Babitha et al., 2008](#) S. Babitha, J.C. Carvahlo, C.R. Soccol, A. Pandey

Effect of light on growth, pigment production and culture morphology of *Monascus purpureus* in solid-state fermentation

World J. Microbiol. Biotechnol., 24 (2008), pp. 2671-2675

[CrossRef](#) ↗ [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Carbonell et al., 2000](#) G.V. Carbonell, H.H. Della, T. Colleta, A.L. Yano, C.E. Darini, Levy B.A.

Fonseca

Clinical relevance and virulence factors of pigmented *Serratia marcescens*

FEMS Immun. Med. Microbiol., 28 (2) (2000), pp. 143-149,

[10.1111/j.1574-695X.2000.tb01469.x](#) ↗

 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Chen et al., 2015](#) W.C. Chen, R.S. Juang, Y.H. Wei

Applications of a lipopeptide biosurfactant, surfactin, produced by microorganisms

Biochem. Eng. J., 103 (2015), pp. 158-169, [10.1016/j.bej.2015.07.009](#) ↗

 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Chen et al., 2016](#) D. Chen, C. Xue, M. Chen, S. Wu, Z. Li

Effects of blue light on pigment biosynthesis of *Monascus*

J. Microbiol., 54 (2016), pp. 305-310

[CrossRef](#) ↗ [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Corrochano, 2007](#) L.M. Corrochano

Fungal photoreceptors: sensory molecules for fungal development and behaviour. Photochemical & photobiological sciences

Euro. Photochem. Assoc. Euro. Soc. Photobiol., 6 (2007), pp. 725-736

[CrossRef](#) ↗ [View in Scopus](#) ↗ [Google Scholar](#) ↗

[De Lucca et al., 2012](#) A.J. De Lucca, C. Carter-Wientjes, K.A. Williams, D. Bhatnagar

Blue light (470 nm) effectively inhibits bacterial and fungal growth

Lett. Appl. Microbiol., 55 (2012), pp. 460-466

[CrossRef ↗](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Dieringer et al., 1977](#) S.M. Dieringer, J.T. Singer, J.J. Cooney

Photokilling of *Micrococcus roseus*

Photochem Photobiol., 26 (1977), pp. 393-396

[CrossRef ↗](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Ferreira et al., 2004](#) C.V. Ferreira, C.L. Bos, H.H. Versteeg, G.Z. Justo, N. Durán, M.P.

Peppelenbosch

Molecular mechanism of violacein-mediated human leukemia cell death

Blood, 104 (2004), pp. 1459-1467

[View in Scopus ↗](#) [Google Scholar ↗](#)

[Gallardo-Casas et al., 2010](#) C.A. Gallardo-Casas, E. Cano-Europa, G.E. López-Galindo, V. Blas-

Valdivia, R. Olvera-Ramírez, M. Franco-Colín, R. Ortiz-Butrón

Phycobiliproteins from *Spirulina maxima* and *Pseudanabaena tenuis*

protect against hepatic damage and oxidative stress caused by Hg²⁺

Rev. Mex. Cienc. Farm., 41 (2010), pp. 30-35

[Google Scholar ↗](#)

[Gitte et al., 2015](#) S.J. Gitte, L.A. Victoria, L.B. Joni, G. Jesse, E. Axel, F.B. Kathleen

Antioxidant and anti-inflammatory properties of an aqueous Cyanophyta extract derived from *Arthrospira platensis*: contribution to bioactivities by the non-phycoyanin aqueous fraction

J. Med. Food, 18 (2015), pp. 535-541

[Google Scholar ↗](#)

[Hall et al., 2016](#) S. Hall, C. McDermott, S. Anoopkumar-Dukie, A.J. McFarland, A. Forbes, A.V.

Perkins, A.K. Davey, R. Chess-Williams, M.J. Kiefel, D. Arora

Cellular effects of pyocyanin, a secreted virulence factor of *Pseudomonas aeruginosa*

Toxins, 8 (2016), p. 236, [10.3390/toxins8080236 ↗](#)

[View in Scopus ↗](#) [Google Scholar ↗](#)

[Herrera-Estrella and Horwitz, 2007](#) A. Herrera-Estrella, B. Horwitz

Looking through the eyes of fungi: molecular genetics of photoreception

Mol. Microbiol., 64 (5) (2007), p. 15

[Google Scholar ↗](#)

[Kodach et al., 2006](#) L.L. Kodach, C.L. Bos, N. Durán, M.P. Peppelenbosch, C.V. Ferreira, J.C.

Hardwick

Violacein synergistically increases 5-fluorouracil cytotoxicity, induces apoptosis and inhibits Akt-mediated signal transduction in human colorectal cancer cells

Carcinogenesis, 27 (2006), pp. 508-516

[CrossRef ↗](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Konzen et al., 2006](#) M. Konzen, D. De Marco, C.A. Cordova, T.O. Vieira, R.V. Antônio, T.B.

Creczynski-Pasa

Antioxidant properties of violacein: possible relation on its biological function

Bioorg Med Chem., 14 (2006), pp. 8307-8313

 [View PDF](#) [View article](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Li and Mira, 2010](#) E. Li, de O.R. Mira

A rapid method for the determination of microbial biomass by dry weight using a moisture analyser with an infrared heating source and an analytical balance

Lett. Appl. Microbiol., 50 (3) (2010), pp. 283-288, [10.1111/j.1472-765X.2009.02789.x ↗](#)

[View in Scopus ↗](#) [Google Scholar ↗](#)

[Liu and Nizet, 2009](#) G.Y. Liu, V. Nizet

Color me bad

Trends Microbiol., 17 (9) (2009), pp. 406-413, [10.1016/j.tim.2009.06.006 ↗](#)

 [View PDF](#) [View article](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Matz et al., 2004](#) C. Matz, P. Deines, J. Boenigk, H. Arndt, L. Eberl, S. Kjelleberg, K. Jürgens

Impact of violacein-producing bacteria on survival and feeding of bacterivorous nanoflagellates

Appl. Environ. Microbiol., 70 (2004), pp. 1593-1599

[View in Scopus ↗](#) [Google Scholar ↗](#)

[Miyake et al., 2005](#) T. Miyake, A. Mori, T. Kii, T. Okuno, Y. Usui

Light effects on cell development and secondary metabolism in *Monascus*

J. Ind. Microbiol. Biotechnol., 32 (2005), pp. 103-108

[CrossRef ↗](#) [View in Scopus ↗](#) [Google Scholar ↗](#)

[Nakamura et al., 2003](#) Y. Nakamura, T. Sawada, Y. Morita, E. Tamiya

Isolation of a psychrotrophic bacterium from the organic residue of a water tank keeping rainbow trout and antibacterial effect of violet pigment produced from the strain

Biochem. Eng., J.12 (2003), pp. 79-86

[CrossRef](#) [Google Scholar](#)

[Nobutaka et al., 2004](#) S. Nobutaka, N. Masami, H. Hiroshi, Y. Isamu, A. Katsumi

Effects of light conditions on prodigiosin stability in the biocontrol bacterium *Serratia marcescens* strain B2

J. Gen. Plant Path., 6 (2004), pp. 367-370

[Google Scholar](#)

[Nora et al., 2017](#) M.E. Nora, S.Y. Aymen, N.E. Hala, A.A. Magdy

Optimization of prodigiosin production by *Serratia marcescens* using crude glycerol and enhancing production using gamma radiation

Biotechnol. Rep., 14 (2017), pp. 47-53, [10.1016/j.btre.2017.04.001](#)

[Google Scholar](#)

[Oren, 2010](#) Aharon Oren

Industrial and environmental applications of halophilic microorganisms

Environ. Technol., 31 (8–9) (2010), pp. 825-834, [10.1080/09593330903370026](#)

[View in Scopus](#) [Google Scholar](#)

[Ortiz-Castro et al., 2014](#) R. Ortiz-Castro, R. Pelagio-Flores, A. Méndez-Bravo, L.F. Ruiz-Herrera, J. Campos-García, J. López-Bucio

Pyocyanin, a virulence factor produced by *Pseudomonas aeruginosa*, alters root development through reactive oxygen species and ethylene signaling in Arabidopsis

Mol. Plant Microbe Interact., 27 (4) (2014), pp. 364-378, [10.1094/MPMI-08-13-0219-R](#)

[View in Scopus](#) [Google Scholar](#)

[Propst and Lubin, 1979](#) C. Propst, L. Lubin

Light-mediated changes in pigmentation of *Pseudomonas aeruginosa* cultures

J. Gen. Microbiol., 113 (1979), pp. 261-266

[CrossRef](#) [View in Scopus](#) [Google Scholar](#)

[Rau and Mitzka-Schnabel, 1985](#) W. Rau, U. Mitzka-Schnabel

Carotenoid synthesis in *Neurospora crassa*

MethodsEnzymol., 110 (1985), pp. 253-267, [10.1016/S0076-6879\(85\)10082-0](#)

 [View PDF](#) [View article](#) [View in Scopus](#) [Google Scholar](#)

[Rebecca et al., 2016](#) S.T. Rebecca, D.S. Vinaya, S.O. Radhika, K.D. Sunil, S.S. Yogesh

Biology and survival of extremely halophilic archaeon *Haloarcula marismortui* RR12 isolated from Mumbai salterns, India in response to

salinity stress

Sci. Rep., 6 (2016), p. 25642, [10.1038/srep25642](https://doi.org/10.1038/srep25642) ↗

[Google Scholar](#) ↗

[Salih et al., 2000](#) A. Salih, A. Larkum, G. Cox, M. Kuhl, Hoegh-Guldberg

Fluorescent pigments in corals are photoprotective

Nature, 408 (2000), pp. 850-853

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[Sánchez et al., 2006](#) C. Sánchez, A.F. Brana, C. Méndez, J.A. Salas

Reevaluation of the violacein biosynthetic pathway and its relationship to indolocarbazole biosynthesis

Chem Bio Chem., 7 (2006), pp. 1231-1240

[CrossRef](#) ↗ [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Seagle et al., 2005](#) B.L. Seagle, K.A. Rezai, Y. Kobori, E.M. Gasyna, K.A. Rezaei, J.R. Norris

Melanin photoprotection in the human retinal pigment epithelium and its correlation with light-induced cell apoptosis

Proc. Natl. Acad. Sci. U.S.A. (2005), pp. 8978-8983

[CrossRef](#) ↗ [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Soni et al., 2009](#) B. Soni, N.P. Visavadiya, D. Madamwar

Attenuation of diabetic complications by C-phycoerythrin in rats: antioxidant activity of C- phycoerythrin including copper -induced lipoprotein and serum oxidation

Br. J. Nutr., 102 (2009), pp. 102-109

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[Soumya et al., 2014](#) K. Soumya, L. Swathi, G.L. Sreelatha, T. Sharmila

Light influences pigment, biomass and morphology in *Chaetomium cupreum* – SS02 – a photo response study

Int. J. Curr. Microbiol. App. Sci., 3 (2014), pp. 53-64

[Google Scholar](#) ↗

[Velmurugan et al., 2010](#) P. Velmurugan, Y.H. Lee, C.K. Venil, P. Lakshmanaperumalsam, J.C. Chae, B. Taek

Effect of light on growth, intracellular and extracellular pigment production by five pigment-producing filamentous fungi in synthetic medium

J. Biosci Bioeng. (2010), pp. 346-350

 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗

[Xu et al., 2014](#) H. Xu, M. Xu, T. Yang, Z. Rao

Effect of temperature on prodigiosin synthesis in *Serratia marcescens*

Chinese, 54 (5) (2014), pp. 517-524

[View in Scopus ↗](#) [Google Scholar ↗](#)

[Yasumasa et al., 1975](#) K. Yasumasa, K. Fumio, O. Shigenobu, T. Takeshi, Y. Saburo

Effect of light on nonphotosynthetic microorganisms

Jpn. J. Microbiol., 19 (5) (1975), pp. 387-393

[Google Scholar ↗](#)

[Yong and Lee, 1991](#) Y.Y.R. Yong, Y.K. Lee

Do carotenoids play a photoprotective role in the cytoplasm of *Haematococcus lacustris* (Chlorophyta)?

Phycologia, 30 (1991), pp. 257-261

[CrossRef ↗](#) [Google Scholar ↗](#)

Cited by (14)

[Screening for pigment production and characterization of pigment profile and photostability in cold-adapted Antarctic bacteria using FT-Raman spectroscopy](#)

2024, Journal of Photochemistry and Photobiology A: Chemistry

[Show abstract](#) ✓

[Isolation and identification of *Methylobacterium komagatae* and its application in textile industries](#)

2024, Natural Product Research

[Show abstract](#) ✓

[Reduction in Pathogenic Biofilms by the Photoactive Composite of Bacterial Cellulose and Nanochitosan Dots under Blue and Green Light ↗](#)

2024, Journal of Functional Biomaterials

[Levan Production by *Suhomyces kilbournensis* Using Sugarcane Molasses as a Carbon Source in Submerged Fermentation ↗](#)

2024, Molecules

Overproduction of Xanthophyll Pigment in *Flavobacterium* sp. JSWR-1 under Optimized Culture Conditions ↗

2024, Journal of Microbiology and Biotechnology

The distribution and drivers of microbial pigments in the cryoconite of four Tibetan glaciers ↗

2024, Environmental Microbiology



[View all citing articles on Scopus ↗](#)

Peer review under responsibility of King Saud University.

© 2019 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University.



All content on this site: Copyright © 2024 Elsevier B.V., its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply.

